

Particle interactions, masses and the symmetry breaking

Fizuli Mamedov

Institute of Physics, Baku, Azerbaijan

Abstract

A possible original $SU(2)_L \times SU(2)_R$ symmetry of the elementary particles and the mechanism of its breaking is discussed. It is concluded that it is the broken symmetry states of the particles which induce the interactions among the particles.

We briefly review the unbroken $SU(2)_L \times U(1)_Y$ symmetry part of the Standard Model of electroweak interactions first for the sake of clarity of the further discussions. According to this theory, the original Lagrangian of the (massless) particles possess both $SU(2)$ and $U(1)_Y$ symmetries for the left - handed fermion doublets. The right - handed fermions have $U(1)_Y$ symmetry only. The triplet of the massless vector gauge bosons associated with the $SU(2)_L$ is $\mathbf{W}_\mu$ and the massless vector gauge boson associated with the $U(1)_Y$ is B_μ . The Lagrangian for the unbroken $SU_L \times U(1)_Y$ symmetry is

$$L = -\frac{1}{4}\mathbf{W}^{\mu\nu}\mathbf{W}_{\mu\nu} - \frac{1}{4}B^{\mu\nu}B_{\mu\nu} + \bar{\Psi}i\gamma^\mu D_\mu\Psi \quad (1)$$

with Ψ denoting the fermionic fields.

The covariant derivative $D_\mu = \partial_\mu + ig\mathbf{W}_\mu\mathbf{T} + \frac{1}{2}ig' B_\mu Y$. $\mathbf{T}$ denotes the Pauli matrices and Y is the hypercharge of the particles. The second term in D_μ act only on the left - handed fermions. The Lagrangian is invariant under the infinitesimal local gauge transformations for $SU(2)_L$ and $U(1)_Y$:

$SU(2)_L$

$$\begin{aligned} \Psi_L &\rightarrow [1 - igT\alpha(x)]\Psi_L & \Psi_R &\rightarrow \Psi_R \\ W_\mu &\rightarrow W_\mu + \partial_\mu\alpha(x) + g\alpha(x) \times W_\mu & B_\mu &\rightarrow B_\mu \end{aligned} \quad (2)$$

$U(1)_Y$

$$\begin{aligned} \Psi_L &\rightarrow [1 - i\frac{1}{2}g'Y\beta(x)]\Psi_L & \Psi_R &\rightarrow [1 - i\frac{1}{2}g'Y\beta(x)]\Psi_R \\ W_\mu &\rightarrow W_\mu & B_\mu &\rightarrow B_\mu + \partial_\mu\beta(x) \end{aligned} \quad (3)$$

The following definitions allow to write the interaction Lagrangian of the fermions and gauge bosons in a compact form: $WT = W^+T^+ + W^-T^- + W_3T_3$, where $W_\mu^\pm = \frac{1}{\sqrt{2}}(W_{1\mu} \mp iW_{2\mu})$ and $T^\pm = \frac{1}{\sqrt{2}}(T_1 \pm iT_2)$. The electromagnetic field A_μ should be contained in the $igW_{3\mu}T_3 + ig'\frac{1}{2}B_\mu Y$, in the neutral term of the covariant derivative. By defining $A = W_3\sin\theta_w + B\cos\theta_w$ and $Z = W_3\cos\theta_w - B\sin\theta_w$, this term can be written as

$$igW_3T_3 + ig'\frac{1}{2}BY = iA[g\sin\theta_w T_3 + g'\cos\theta_w \frac{1}{2}Y] + iZ[g\cos\theta_w T_3 - g'\sin\theta_w \frac{1}{2}Y]$$

which yields the quasi - Gell - Mann -Nishijima formula for the electric charge: $ieQ = ie(T_3 + \frac{1}{2}Y)$. For the left - handed fermions $T_3 = \pm\frac{1}{2}$ and for the right - handed fermions $T_3 = 0$. Also $g = e/\sin\theta_w$, $g' = e/\cos\theta_w$, $g_Z = e/(\sin\theta_w\cos\theta_w)$, $x_w = \sin^2\theta_w$.

In terms of the $W_\mu^\pm$, A_μ and Z_μ fields the fermion - gauge boson interaction Lagrangian is written as

$$-L' = eJ_{em}^\mu A_\mu + \frac{g}{\sqrt{2}}(J_L^{+\mu}W_\mu^+ + J_L^{-\mu}W_\mu^-) + g_Z J_Z^\mu Z_\mu \quad (4)$$

Here $J_L^{\pm\mu} = \sqrt{2}\bar{\Psi}\gamma^\mu T_L^\pm\Psi$, $J_Z^\mu = \bar{\Psi}\gamma^\mu[T_{3L} - x_w Q]\Psi$ and $J_{em}^\mu = \bar{\Psi}\gamma^\mu Q\Psi$ are the fermion - gauge field interaction currents (T_L acts on the left - handed fermions only). The Standard Model scenario spontaneous symmetry breaking does not affect this form of the interaction currents.

It is possible that the symmetry breaking occurs because of the fusion and condensation of the certain type of the fermions. In such a picture only massless fermionic fields exist at the initial, highest symmetry stage. Left - handed and right - handed fermions manifest themselves as two separate irreducible representations of the most fundamental field. Besides the space-time spin s , fermions also have the electroweak spin¹ $I_W = \frac{1}{2}$ for both left - handed and right - handed fermions. The fused and condensed states of the left - handed fermions can be separated into electroweak spin triplet $I_W = 1$ and singlet $I_W = 0$ states.

$$|s_L = \frac{1}{2} > |I_W = \frac{1}{2}, m_W = \pm\frac{1}{2} > \times |s_L = \frac{1}{2} > |I_W = \frac{1}{2}, m_W = \pm\frac{1}{2} > =$$

$$|s_L = 1 > (|I_W = 1 > + |I_W = 0 >) \quad (5)$$

The electroweak triplet states are represented by the triplet of the vector gauge boson fields² $(\mathbf{W}^\mu)_L$ and the singlet state is represented by the vector boson field $(B^\mu)_L$. The similar fusion and condensation occurs for the right - handed fermions.

$$|s_R = \frac{1}{2} > |I_W = \frac{1}{2}, m_W = \pm\frac{1}{2} > \times |s_R = \frac{1}{2} > |I_W = \frac{1}{2}, m_W = \pm\frac{1}{2} > =$$

$$|s_R = 1 > (|I_W = 1 > + |I_W = 0 >) \quad (6)$$

The states with $s_L = 1$, $(\mathbf{W}^\mu)_L$ and $(B^\mu)_L$ couple to the left-handed fermions only and the states with $s_R = 1$, $(\mathbf{W}^\mu)_R$ and $(B^\mu)_R$ couple to the right - handed fermions only.

¹We use the usual weak isospinor notation, I_W for the electroweak spin in this work

²It is the space - time parts of these fields that are treated as classical fields at first and after the second quantization become quantum fields

The states $\frac{1}{\sqrt{2}}((W_1^\mu)_L + i(W_2^\mu)_L)$, $\frac{1}{\sqrt{2}}((W_1^\mu)_L - i(W_2^\mu)_L)$ and $(W_3^\mu)_L$ correspond to the three generators of the irreducible representation of the $SU(2)_L$ ($I_W = 1$) in the spherical coordinates. Therefore the fields $\frac{1}{\sqrt{2}}((W_1^\mu)_L \pm i(W_2^\mu)_L)$ should be associated with the $I_W = 1$, $m_W = \pm 1$ states, and the field W_3^μ with the $I_W = 1$, $m_W = 0$ state. Thus all the particles (including the fermions) correspond to the combinations of the fields with the definite value of m_W (the role of the m_W is played by T_3 in the context of the $SU(2)_L \times U(1)_Y$ symmetry, discussed earlier in this work). Since both $(W_3^\mu)_L$ and $(B^\mu)_L$ have $m_W = 0$, their mixtures will also have the definite value for the m_W , $m_W = 0$, and can exist as separate particles. The (massive) Z boson³ and the (massless) photon are indeed two such particles. Nature separates the 'steam' part of the $m_W = 0$ mixed states, the massless vector boson from their condensate part, the massive vector boson. Interestingly, the fields with $m_W = 0$, the Z bosons and photons carry the interactions between the same type of fermions, the particles of the same handedness, m_W and generation. On the other hand, fields with $|m_W| = 1$, the $W_\mu^\pm$, couple fermions with the different m_W only and can couple particles of the different generations.

Let us assume that the $\mathbf{W}^\mu$ gain the mass M_W and the B^μ gain the mass M_B as a result of the symmetry breaking. It is natural to assume that, due to the universality of the condensation and couplings,

$$M_W/g = M_B/g' \quad (7)$$

This equation implies that the masses gained by the $\mathbf{W}$ and B due to the symmetry breaking are proportional to their respective coupling strengths, $M_W = kg$ and $M_B = k'g'$, with $k = k'$. Eq.(7) leads to the usual definition of Z_μ and to the well - known $M_W - M_Z$ mass relation:

$$\begin{aligned} M_W^2 W_1^2 + M_W^2 W_2^2 + M_W^2 W_3^2 + M_B^2 B^2 = \\ M_W^2 W^+ W^- + (M_W^2/g^2)(g^2 W_3^2 + (g')^2 B^2) = \\ M_W^2 W^+ W^- + (M_W^2/g^2)(gW_3 - g'B)^2 + 4(M_W^2/g^2)gg'W_3B = \\ M_W^2 W^+ W^- + M_Z^2 Z^2 \end{aligned}$$

with $M_Z = M_W \sqrt{g^2 + (g')^2}/g = M_W/\cos\theta_W$ and

$Z = (gW_3 - g'B)/\sqrt{g^2 + (g')^2} = W_3 \cos\theta_W - B \sin\theta_W$. The term $4(M_W^2/g^2)gg'W_3B$ in this equation disappears because W_3 and B are orthogonal: these states belong to the different multiplets of the electroweak spin I_W .

The known values for the electroweak spin and hypercharge of the particles leads to the following possible scenario of the symmetry breaking process within this model. At first only B_μ fields couple to the fermions. The $(B^\mu)_L$ to the left - handed fermions and the $(B^\mu)_R$ to the right - handed fermions only. The

³We will often drop the indices L and R when their presence is not important

fermion doublets have $SU(2)_L \times SU(2)_R$ global symmetry at this stage with the electroweak quantum numbers:

$$\nu_L \quad m_W = \frac{1}{2}, \quad Y = -1 \qquad e_L \quad m_W = -\frac{1}{2}, \quad Y = -1$$

$$u_L \quad m_W = \frac{1}{2}, \quad Y = \frac{1}{3} \qquad d_L \quad m_W = -\frac{1}{2}, \quad Y = \frac{1}{3}$$

$$\nu_R \quad m_W = \frac{1}{2}, \quad Y = -1 \qquad e_R \quad m_W = -\frac{1}{2}, \quad Y = -1$$

$$u_R \quad m_W = \frac{1}{2}, \quad Y = \frac{1}{3} \qquad d_R \quad m_W = -\frac{1}{2}, \quad Y = \frac{1}{3}$$

The $SU(2)_L \times SU(2)_R$ global symmetry breaking occurs when $(B^\mu)_L$ also couples to the right - handed fermions. Simultaneously, the direct left - right handed fermion couplings, the masses of the fermions appear. $(B^\mu)_R$ does not couple to the left - handed fermions: in that case the original $SU(2)_L \times SU(2)_R$ global symmetry would be preserved. Typically the emergence of the interactions (couplings) is accompanied with the movement of the system to the less symmetrical state. For example, the inclusion of the two - particle interactions between the atoms replaces the global $SO(3)$ symmetry by the global $U(1)$ symmetry. Besides, the $(B^\mu)_L$ couples to the upper members of the right - handed doublets with the strength $Y = 1$ and to the lower members of these doublets with the strength $Y = -1$ (the $(B^\mu)_L$ couples 'color - blindly', with $|Y| = 1$, to the quarks of the different irreducible representation of the most fundamental field, to the u_R and d_R). Once again, the same $(B^\mu)_L$ couplings to these particles would preserve the right - handed doublets and hence the global $SU(2)_L \times SU(2)_R$ symmetry. As a result, the well - known values of the electroweak quantum numbers of the particles are produced:

$$\nu_L \quad m_W = \frac{1}{2}, \quad Y = -1 \qquad e_L \quad m_W = -\frac{1}{2}, \quad Y = -1$$

$$u_L \quad m_W = \frac{1}{2}, \quad Y = \frac{1}{3} \qquad d_L \quad m_W = -\frac{1}{2}, \quad Y = \frac{1}{3}$$

$$\nu_R \quad m_W = 0, \quad Y = 0 \qquad e_R \quad m_W = 0, \quad Y = -2$$

$$u_R \quad m_W = 0, \quad Y = \frac{4}{3} \qquad d_R \quad m_W = 0, \quad Y = -\frac{2}{3}$$

The $(\mathbf{W}^\mu)_L$ triplet of the vector bosons' coupling to the left - handed fermions also occurs at this stage. From Eq.(7) we see that indeed a non-zero value of g produces a non-zero value for M_W , once the B^μ gains its mass at the stage of its own coupling to the fermions. With no right - handed fermion doublets around, $(W^\mu)_R$ does not couple to the fermions. Also, the coupling of the $(W^\mu)_R$ to the fermions would produce a local $SU(2)_L \times SU(2)_R$ symmetry, not a typical consequence of the appearance of the new interactions. Thus the emergence of the new types of interactions among the particles leads to the disappearance of the global $SU(2)_L \times SU(2)_R$ symmetry, along the way also breaking the space left - right symmetry. The fusion of the fermions and generation of the B^μ mass also breaks the symmetry, the original symmetry of the most fundamental field, the irreducible representations of which are the non - interacting massless fermions. Thus we conclude that the interactions (couplings) among the particles are the manifestations of the symmetry breaking. In other words, it is the broken symmetry states that induce the motion of the particles along the curved paths and the mass, as in the case of the gravitational field, is the measure of this curvedness. It is interesting to notice that the 'unbroken part' of the $W_3^\mu - B^\mu$ mixture, the photons do not interact with the neutrinos, the particles which avoid at least the mass generation consequence of the symmetry breaking.

It is the $s_L = 1$ state with $I_W = 0$, $(B^\mu)_L$, that couples to the right - handed fermions after the $SU(2)_L \times SU(2)_R$ symmetry is broken. Therefore the left - handed fermions and the right - handed fermions might have two distinct electroweak spin quantum numbers, $(I_W)_L = \frac{1}{2}$ and $(I_W)_R = \frac{1}{2}$, respectively.

The fermion - vector boson interaction Lagrangian in the form of Eq.(4) appears only after the breaking of the global $SU(2)_L \times SU(2)_R$ symmetry, after its replacement of by the $U(1)$ electromagnetic gauge symmetry. As for the $SU(2)_L$ gauge symmetry transformations, Eq.(2) and, perhaps, $U(1)_Y$ symmetry transformations, Eq.(3), they are never realized for the electroweak interactions Lagrangian within this model.

References

1. S. L. Glashow, *Nuclear Physics* **22** (1961), 579
2. S. Weinberg, *Physical Review Letters*, **19**, 1264
3. A. Salam, *Svartholm, N, Proceedings of the Eighth Nobel Symposium, Almqvist & Wiksell, Stockholm* (1968)
4. V. Barger, R. Philips, *Collider Physics, Addison - Wesley Publishing Company, Inc*, (1997)